

HDL-TR-1897

February 1980

AD A 08 1982

Assessment of $\text{Pr}^{3+}:\text{KY}_3\text{F}_{10}$ as a Blue-Green Laser

by Clyde A. Morrison
Donald E. Wortman
Richard P. Leavitt
Hans P. Jenssen

DTIC
ELECTE
APR 11 1980
S A D



U.S. Army Electronics Research
and Development Command
Harry Diamond Laboratories

Adelphi, MD 20783

Approved for public release; distribution unlimited.

80 4 11 001

DDC FILE COPY

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

Citation of manufacturers' or trade names does not constitute an official indorsement or approval of the use thereof.

Destroy this report when it is no longer needed. Do not return it to the originator.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER HDL-TR-1897	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Assessment of Pr ³⁺ :KY ₃ F ₁₀ as a Blue-Green Laser.	5. TYPE OF REPORT & PERIOD COVERED Technical Report	
6. PERFORMING ORG. REPORT NUMBER		7. AUTHOR(s) Clyde A./Morrison Hans P./Jenssen Donald P./Wortman (Massachusetts Richard P./Leavitt Inst. of Tech.)
8. CONTRACT OR GRANT NUMBER(s)		9. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Program Ele: 61.15.3N
10. CONTROLLING OFFICE NAME AND ADDRESS Harry Diamond Laboratories 2800 Powder Mill Road Adelphi, MD 20783		11. REPORT DATE February 1980
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 27
14. SECURITY CLASS. (of this report) Unclassified		15. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES HDL Project: 355832 DRCMS Code: 36AA.600061153		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Rare earth Optical spectra Praseodymium Optical transition probabilities KY ₃ F ₁₀ Rare-earth laser		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The reported energy levels of praseodymium and europium in the host material KY ₃ F ₁₀ have been analyzed. Using the x-ray data, lattice sums have been done, and odd crystal-field parameters have been calculated. Using these results, the transition probabilities for both magnetic and electric dipoles have been calculated for the rare earth ions. Discussions are given of the application of the results to a blue-green laser using praseodymium.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

CONTENTS

	<i>Page</i>
1. INTRODUCTION	5
2. CRYSTALLOGRAPHIC DATA AND LATTICE SUM FOR KY_3F_{10}	6
3. ANALYSIS OF Eu^{3+} , Nd^{3+} , AND Pr^{3+} OPTICAL DATA	7
4. CALCULATIONS AND DISCUSSION OF RESULTS	8
LITERATURE CITED	22

TABLES

1. Crystal Structure Data on KY_3F_{10}	7
2. Lattice Sum for KY_3F_{10}	7
3. Phenomenological Crystal Field Parameters (B_{kq}) for Triply Ionized Lanthanides in KY_3F_{10}	8
4. Even-K Phenomenological A_{kq} and A_{kq} Determined from Summation over KY_3F_{10} Lattice	9
5. Crystal Field Parameters for Triply Ionized Lanthanides in KY_3F_{10}	9
6. Energy Levels for Ce^{3+} in KY_3F_{10}	9
7. Energy Levels for Pr^{3+} in KY_3F_{10}	10
8. Energy Levels for Nd^{3+} in KY_3F_{10}	11
9. Energy Levels for Pm^{3+} in KY_3F_{10}	12
10. Energy Levels for Sm^{3+} in KY_3F_{10}	13
11. Energy Levels for Eu^{3+} in KY_3F_{10}	14
12. Energy Levels for Gd^{3+} in KY_3F_{10}	15
13. Energy Levels for Tb^{3+} in KY_3F_{10}	16
14. Energy Levels for Dy^{3+} in KY_3F_{10}	17
15. Energy Levels for Ho^{3+} in KY_3F_{10}	18
16. Energy Levels for Er^{3+} in KY_3F_{10}	19

17.	Energy Levels for Tm^{3+} in KY_3F_{10}	20
18.	Energy Levels for Yb^{3+} in KY_3F_{10}	21
19.	Branching Ratios from $^3\text{P}_0$ Energy Level to Lower Lying J-Multiplets for Pr^{3+} in KY_3F_{10} and in YLF	21
20.	Electric Dipole Transition Probabilities from Fluorescing $^3\text{P}_0$ Energy Level to $^3\text{H}_4$ Energy Levels for Pr^{3+} in KY_3F_{10} and in YLF	21

1. INTRODUCTION

The most promising blue-green laser crystal reported¹ to date is $\text{Pr}^{3+}:\text{YLF}$ (praseodymium in lithium yttrium fluoride, LiYF_4 or YLF). The only strong transitions from the $^3\text{P}_0$ fluorescing level to the $^3\text{H}_4$ ground state multiplet of Pr^{3+} are, unfortunately, to the lowest lying levels of $^3\text{H}_4$, which means that the laser is a three-level system even at low temperature (77 K). Since four-level laser action is necessary to achieve low lasing threshold and high efficiency, other hosts where the laser terminates on an upper energy level are desired, providing that the material has other required properties for blue-green laser operation such as a large band gap and good optical quality. One possible material is $\text{Pr}^{3+}:\text{KY}_3\text{F}_{10}$, which was examined in this work as described below.

X-ray data² were used in a lattice sum calculation, as is discussed in section 2, to determine the spherical tensor amplitudes, A_{kq} , of the electrostatic field at the impurity ion site. By using these A_{kq} in a three-parameter model³ of crystal fields, it was possible to estimate the energy levels of europium, Eu^{3+} , in KY_3F_{10} , which were obtained from the optical data recorded⁴ by Porcher and Caro for $\text{Eu}^{3+}:\text{KY}_3\text{F}_{10}$ powder. These data were then successfully analyzed, and phenomenological crystal field parameters, B_{kq} , were found by a method described elsewhere⁵ that gave a least-squares fit of 6.9 cm^{-1} between 26 calculated and measured energy levels for Eu^{3+} .

The above Eu^{3+} parameters were scaled⁶ by the appropriate ρ_k values, $B_{kq} = \rho_k$ (ion dependent) A_{kq} (host dependent), to give even-fold (even- k) B_{kq} required to predict the Stark split energy levels for neodymium, Nd^{3+} , and Pr^{3+} ; the method of calculation is described elsewhere.⁷ One of the authors had available preliminary (and unpublished) energy levels and optical data for both $\text{Nd}^{3+}:\text{KY}_3\text{F}_{10}$ and $\text{Pr}^{3+}:\text{KY}_3\text{F}_{10}$. These data were analyzed, and phenomenological B_{kq} were found that gave root mean square (rms) deviations of 6.3 cm^{-1} and 14.3 cm^{-1} , respectively, between 23 Nd^{3+} and 35 Pr^{3+} calculated and measured energy levels.

By using the even- k phenomenological B_{kq} for Pr^{3+} and the odd- k A_{kq} obtained from the lattice sum, electric and magnetic dipole transition probabilities and branching ratios were calculated. According to the branching ratio calculations, about 33 percent of the electromagnetic radiation from the $^3\text{P}_0$ level goes to the ground multiplet, which compares⁷ with 36 percent going to the ground multiplet for Pr^{3+} in YLF. The largest transition goes to the lower energy states, as for Pr^{3+} in YLF. However, two of the largest three transitions from $^3\text{P}_0$ in KY_3F_{10} go to levels at 166 and 217 cm^{-1} , whereas one of the larger transitions goes to a level at 79 cm^{-1} in YLF. Thus, four-level operation of Pr^{3+} in KY_3F_{10} at 77 K might be attained since the possible terminal level would not be so thermally populated as it is in YLF.

¹ L. Esterowitz, R. Allen, M. Kruer, F. Bartoli, L. S. Goldberg, H. P. Jenssen, A. Linz, and V. Nicolai, *J. Appl. Phys.*, **48** (1977), 650.

² J. W. Pierce and H. Y.-P. Hong, *Proceedings of Tenth Rare-Earth Research Conference* (1973), 527.

³ Richard P. Leavitt, Clyde A. Morrison, and Donald E. Wortman, *Rare-Earth Ion-Host Crystal Interactions 3. Three-Parameter Theory of Crystal Fields*, Harry Diamond Laboratories HDL-TR-1673 (June 1975).

⁴ P. Porcher and P. Caro, *J. Chem. Phys.*, **65** (1976), 89.

⁵ N. Karayianis, D. E. Wortman, and H. P. Jenssen, *J. Phys. Chem. Solids*, **37** (1976), 675.

⁶ Clyde A. Morrison, Nick Karayianis, and Donald E. Wortman, *Rare-Earth Ion-Host Lattice Interactions 4. Predicting Spectra and Intensities of Lanthanides in Crystals*, Harry Diamond Laboratories HDL-TR-1816 (June 1977).

⁷ Leon Esterowitz, Filbert J. Bartoli, Roger E. Allen, Donald E. Wortman, Clyde A. Morrison, and Richard P. Leavitt, *Energy Levels and Line Intensities for the Ground Configuration of Pr^{3+} in LiYF_4* , Harry Diamond Laboratories HDL-TR-1875 (February 1979).

In addition to the above calculations, phenomenological A_{kq} were determined by the relation $B_{kq} = \rho_k A_{kq}$ and from the phenomenological B_{kq} determined in the Pr^{3+} , Nd^{3+} , and Eu^{3+} spectral analyses. These A_{kq} , which should depend only upon the host crystal, were then averaged to give a set of A_{kq} at a Y site. This set of A_{kq} were multiplied by appropriate ρ_k for each lanthanide ion, and B_{kq} were obtained for the entire lanthanide series. Energy levels for each of the triply ionized lanthanides in KY_3F_{10} were calculated by using the appropriate set of B_{kq} , and these results are given.

2. CRYSTALLOGRAPHIC DATA AND LATTICE SUM FOR KY_3F_{10}

The first complete x-ray analysis of KY_3F_{10} was done by Pierce and Hong.² The space group most consistent with their x-ray data is $\text{Fm}\bar{3}\text{m}$ (space group 225 of the international tables³) with all the Y ions at sites of C_{4v} symmetry. A second possible space group, $\text{F}\bar{4}3\text{m}$, fit Pierce and Hong's data as well as the $\text{Fm}\bar{3}\text{m}$, but the $\text{Fm}\bar{3}\text{m}$ was chosen by them because it is more symmetric than $\text{F}\bar{4}3\text{m}$. This choice has been confirmed by recent optical data⁴ on Eu^{3+} in KY_3F_{10} . If the space group were $\text{F}\bar{4}3\text{m}$, the Eu^{3+} ion would occupy a site of C_{2v} symmetry, and complete removal of the degeneracy of the free ion would be observed. However, the optical spectra agree with what should be observed if the ion were in a site of C_{4v} symmetry, which agrees with the symmetry of the Y site of the space group $\text{Fm}\bar{3}\text{m}$.

Recently, the material potassium ytterbium fluoride ($\text{KYb}_3\text{F}_{10}$) has been grown⁹ and found to have the same space group, $\text{Fm}\bar{3}\text{m}$, with the cubic cell dimension, $a = 11.431 \text{ \AA}$, and the positions of the constituent ions are $x_{\text{Yb}} = 0.24$, $x_{\text{F}_1} = 0.33$, and $x_{\text{F}_2} = 0.12$. A large number of compounds also have been grown recently by Vedrine et al.,¹⁰ which are cubic and have the form $\text{KLn}_3\text{F}_{10}$ ($\text{Ln} = \text{lutetium—Lu, Yb, thulium—Tm, erbium—Er, holmium—Ho, or dysprosium—Dy}$) and $\text{RbLn}_3\text{F}_{10}$ ($\text{Ln} = \text{terbium—Tb, gadolinium—Gd, Eu, or samarium—Sm}$) with only the cubic cell dimension reported.

We have performed the lattice sum for KY_3F_{10} with the origin at a Y site using Pierce and Hong's² data as given in table 1 with the various ions occupying the sites given in the international tables,³ and the results are given in table 2. There are 24 Y ions in a unit cell, and each site has C_{4v} symmetry. However, eight Y ions at $(0, 0, \pm x)$ have the principal axis of C_{4v} along z , eight Y ions at $(\pm x, 0, 0)$ have the principal axis along x , and eight Y ions at $(0, \pm x, 0)$ have the principal axis along y . The full 24 Y ions are generated by three additional translations within the unit cell: $(0, \frac{1}{2}, \frac{1}{2})$, $(\frac{1}{2}, 0, \frac{1}{2})$, and $(\frac{1}{2}, \frac{1}{2}, 0)$, which preserve the axis of symmetry of each particular site. Whereas the optical spectra of a rare-earth ion at a single site possessing C_{4v} symmetry have well-defined polarization selection rules, the multiplicity of the principal axes causes polarization data to be meaningless. Because of the multiplicity of the principal axes of C_{4v} symmetry in KY_3F_{10} , a magnetic field applied in an arbitrary direction gives rise to three sets of Zeeman splittings in the spectrum of a rare-earth ion. These sets of lines interchange as the direction of the magnetic field is varied. It is possible that the only advantage of using a magnetic field in analyzing the optical spectra would be for those rare-earth ions possessing an even number of electrons. The magnetic field could then be used to readily identify the doublets existing for these ions in C_{4v} symmetry.

² J. W. Pierce and H. Y.-P. Hong, Proceedings of Tenth Rare-Earth Research Conference (1973), 527.

³ P. Porcher and P. Caro, J. Chem. Phys., 65 (1976), 89.

⁴ International Tables, I, Kynoch Press, Birmingham, U.K. (1952), 338, table 225.

⁹ M. Labeau, S. Aleonard, A. Vedrine, R. Boutonnet, and J. C. Cousseins, Mat. Res. Bull., 9 (1974), 615.

¹⁰ A. Vedrine, R. Boutonnet, R. Sabatier, and J. Cousseins, Bull. Soc. Chim. Fr., 3-4 (1975), 445.

TABLE 1. CRYSTAL STRUCTURE DATA ON KY_3F_{10}
Space Group $Fm\bar{3}m$ cubic $a = 11.542 \pm 0.004 \text{ \AA}$

Ion	Position	Site symmetry	Ion ^a position
Y	24e	C_{4v}	$x = 0.2401$
K	8c	T_d	—
F ₁	48i	C_{2v}	$x = 0.3353$
F ₂	32f	C_{3v}	$x = 0.1081$

^a These are the ion positions that have to be determined by x-ray analysis for use in the International Tables. The position 8c is determined in terms of rational fractions.

Sources: J. W. Pierce and H. Y.-P. Hong, Proceedings of Tenth Rare-Earth Research Conference (1973), 527; International Tables, 1, Kynoch Press, Birmingham, U.K. (1952), 338, table 225.

TABLE 2. LATTICE SUM FOR KY_3F_{10}

Spherical tensor amplitudes	
A_{kq}	($\text{cm}^{-1}/(\text{\AA})^k$)
A_{10}	4749.97
A_{20}	-4011.53
A_{30}	-4431.00
A_{40}	-4822.27
A_{44}	1531.83
A_{50}	1048.16
A_{54}	2799.43
A_{60}	661.89
A_{64}	42.32
A_{70}	54.69
A_{74}	241.86

Note: The origin is the yttrium ion at (0, 0, $\pm x$).
The effective charges are the normal valence values.

Source of x-ray data: J. W. Pierce and H. Y.-P. Hong, Proceedings of Tenth Rare-Earth Research Conference (1973), 527.

3. ANALYSIS OF Eu^{3+} , Nd^{3+} , AND Pr^{3+} OPTICAL DATA

In making the energy level calculations, the crystal field Hamiltonian,

$$H_x = \sum_{kq} B_{kq} C_{kq}, \quad (1)$$

where C_{kq} are spherical tensors, was diagonalized in a basis of states spanning a particular set of free-ion J-multiplets by the method described elsewhere.⁵ The B_{kq} are related to the A_{kq} , which were obtained in the lattice sum, by the expression,

$$B_{kq} = \rho_k A_{kq}. \quad (2)$$

The ρ_k were obtained^{3,6} by considering shielding and radial wave-function expansion and are ion-dependent quantities given by Morrison et al in their table II.⁶

³ Richard P. Leavitt, Clyde A. Morrison, and Donald E. Wortman, Rare-Earth Ion-Host Crystal Interactions 3, Three-Parameter Theory of Crystal Fields, Harry Diamond Laboratories HDL-TR-1673 (June 1975).

⁵ N. Karayianis, D. E. Wortman, and H. P. Jessen, J. Phys. Chem. Solids, 37 (1976), 675.

⁶ Clyde A. Morrison, Nick Karayianis, and Donald E. Wortman, Rare-Earth Ion-Host Lattice Interactions 4, Predicting Spectra and Intensities of Lanthanides in Crystals, Harry Diamond Laboratories HDL-TR-1816 (June 1977).

In this work, the A_{kq} given in table 2 for KY_3F_{10} were multiplied by the ρ_k values⁶ for Eu^{3+} . This procedure gave a set of B_{kq} , which were used to calculate an initial set of energy levels for $Eu^{3+}:KY_3F_{10}$. These calculated levels were compared with the experimental values of Porcher and Caro⁴ in a least-squares fit calculation. The B_{kq} were varied, and new calculated energy levels were determined in the fitting procedure until a least rms value of 6.9 cm^{-1} between 26 calculated and experimental levels was obtained. The phenomenological B_{kq} yielding this fit for Eu^{3+} are given in table 3. Porcher and Caro⁴ reported B_{kq} for Eu^{3+} , and these values are given for comparison in table 3.

TABLE 3. PHENOMENOLOGICAL CRYSTAL FIELD PARAMETERS (B_{kq}) FOR TRIPLY IONIZED LANTHANIDES IN KY_3F_{10}

Parameter	Pr ^a	Nd ^a	Eu ^b	Eu ^c
$B_{20}\text{ (cm}^{-1}\text{)}$	-589.0	-621.1	-517.6	-551
$B_{40}\text{ (cm}^{-1}\text{)}$	-1711	-1566	-1337	-1360
$B_{44}\text{ (cm}^{-1}\text{)}$	518.9	252.2	359.6	345
$B_{60}\text{ (cm}^{-1}\text{)}$	656.9	366.7	400.1	394
$B_{64}\text{ (cm}^{-1}\text{)}$	-195.9	-366.6	-2 [*]	-234
Root mean square	14.3	6.3	6.9	7.6
Levels in fit (No.)	35	23	26	25
Multiplets with data (No.)	12	5	8	8
Crystal field splittings (No.)	23	18	18	16

^a Preliminary data from H. P. Jessen, Massachusetts Institute of Technology (unpublished).

^b Our fit on data reported by P. Porcher and P. Caro, J. Chem. Phys., 65 (1976), 89.

^c B_{64} reported by Porcher and Caro. B_{64} should be -234 rather than +234 (private communication).

The phenomenological B_{kq} for Eu^{3+} that we obtained were scaled by the ratio $\rho_k(Nd)/\rho_k(Eu)$ to give an initial set of B_{kq} for Nd^{3+} . This initial set of B_{kq} was used to calculate energy levels for Nd^{3+} in KY_3F_{10} for comparison with our preliminary data. Phenomenological B_{kq} for Nd^{3+} were obtained that yielded a least rms value of 6.3 cm^{-1} between 23 calculated and measured levels; these B_{kq} are given in table 3.

Initial B_{kq} were obtained for Pr^{3+} in KY_3F_{10} by using the phenomenological B_{kq} for Eu^{3+} . These B_{kq} were used to calculate energy levels that aided us in identifying 35 energy levels in the Pr^{3+} ground configuration. Phenomenological B_{kq} listed in table 3, gave a least rms deviation of 14.3 cm^{-1} between 35 calculated and measured Pr^{3+} energy levels.

4. CALCULATIONS AND DISCUSSION OF RESULTS

The above phenomenological B_{kq} for Pr, Nd, and Eu were divided by the appropriate⁶ ρ_k values, and the phenomenological A_{kq} values listed in table 4 were obtained. Since the A_{kq} are supposed to be host dependent only, the variations in the A_{kq} give some idea about the uncertainty in these parameters. The even-k A_{kq} determined from these three ions were averaged to give a set of A_{kq} for KY_3F_{10} ; these values are given in table 4 along with the A_{kq} obtained in the lattice sums for two different charges on the F ions. In addition, the averaged A_{kq} were multiplied by the appropriate⁶ ρ_k , and B_{kq} were determined for the lanthanides in KY_3F_{10} as given in table 5. Energy levels calculated by using the B_{kq} given in table 5 for each of the lanthanides in KY_3F_{10} are given in tables 6 to 18. In the tables, 2 MU is twice the crystal quantum number,¹¹ 2μ .

⁴ P. Porcher and P. Caro, J. Chem. Phys., 65 (1976), 89.

⁶ Clyde A. Morrison, Nick Karayianis, and Donald E. Wortman, Rare-Earth Ion-Host Lattice Interactions 4. Predicting Spectra and Intensities of Lanthanides in Crystals, Harry Diamond Laboratories HDL-TR-1816 (June 1977).

¹¹ K. H. Hellwege, Ann. Physik, 4 (1948), 95.

TABLE 4. EVEN-K PHENOMENOLOGICAL A_{k0} AND A_{k0} DETERMINED FROM SUMMATION OVER KY_3F_{10} LATTICE

Amplitude or radial factor	Calculated crystal field parameter					
	1	2	3	4	5	6
A_{20}	-3354	-3641	-3107	-3367	-4011	-4288
A_{40}	-2647	-2711	-2765	-2708	-4822	-4752
A_{44}	802.7	436.6	743.6	661.0	1532	1493
A_{60}	350.3	230.7	320.0	300.3	661.9	654.7
A_{64}	-104.5	-230.6	-217.3	-184.1	42.32	42.1
ρ_2	0.1756	0.1706	0.1666	—	—	—
ρ_4	0.6464	0.5776	0.4836	—	—	—
ρ_6	1.8754	1.5897	1.2503	—	—	—

Notes:

1: $A_{k0} = B_{k0}/\rho_k$; $Pr^{3+}B_{k0}$ and ρ_k .

2: $A_{k0} = B_{k0}/\rho_k$; $Nd^{3+}B_{k0}$ and ρ_k .

3: $A_{k0} = B_{k0}/\rho_k$; $Eu^{3+}B_{k0}$ and ρ_k .

4: Average of 1, 2, and 3.

5: A_{k0} from lattice sum where the charges used in the sum are $q_k = 1$, $q_4 = 3$, and $q_6 = -1$.

6: A_{k0} from lattice sum with $q_k = 0.9$, $q_4 = 3$, and $q_6 = -0.99$.

TABLE 5. CRYSTAL FIELD PARAMETERS FOR TRIPLY IONIZED LANTHANIDES IN KY_3F_{10}

Ion	N ^a	N ^b	B_{20}	B_{40}	B_{44}	B_{60}	B_{64}	Root mean square deviation
Ce	0	7	-620	-2041	498	703	-431	—
Pr	35	70	-589	-1711	519	657	-196	14.3
Nd	23	60	-621	-1566	252	367	-367	6.3
Pm	0	89	-565	-1446	353	427	-262	—
Sm	0	59	-562	-1367	334	397	-243	—
Eu	26	59	-518	-1337	360	400	-272	6.9
Gd	0	67	-562	-1261	308	356	-219	—
Tb	0	75	-563	-1216	297	337	-207	—
Dy	0	67	-566	-1176	287	319	-195	—
Ho	0	68	-570	-1142	279	304	-186	—
Er	0	59	-574	-1117	273	295	-181	—
Tm	0	70	-580	-1098	268	290	-178	—
Yb	0	7	-595	-1056	260	274	-168	—

^a Number of experimental levels.

^b Number of theoretical levels.

Note: Units are in cm^{-1} .

TABLE 6. ENERGY LEVELS FOR Ce^{3+} IN KY_3F_{10}

LSJ state (centroid)		Energy level			
$2F \frac{5}{2}$ (250)	2 MU TH.	1 0	3 342	3 600	
$2F \frac{7}{2}$ (2550)	2 MU TH.	1 2271	3 2624	1 2695	3 3074

TABLE 7. ENERGY LEVELS FOR Pr^{3+} IN KY_3F_{10}

LSJ state (centroid)		Energy level							
3H 4 (232)	2 MU	2	4	0	4	2	0	0	
	TH.	-7	69	161	183	223	511	541	
	EXP.	0		166	175	217			
3H 5 (2361)	2 MU	2	0	4	0	2	4	0	2
	TH.	2197	2222	2226	2280	2313	2334	2393	2656
	EXP.	2184		2211					2684
3H 6 + 3F 2 (4546, 5080)	2 MU	0	2	0	4	2	0	4	2
	TH.	4290	4354	4371	4430	4452	4524	4531	4562
	EXP.	4283	4347		4418	4477			
	2 MU	4	4	4	4	2	0		
	TH.	4991	4993	5031	5071	5100	5214		
	EXP.			5041		5075	5229		
3F 3 (6466)	2 MU	2	4	4	2	0			
	TH.	6422	6450	6458	6499	6573			
	EXP.	6433	6441		6496	6573			
3F 4 (6928)	2 MU	2	4	4	0	0	2	0	
	TH.	6801	6804	6909	6994	7006	7068	7150	
	EXP.	6829		6895		7006	7055		
1G 4 (9945)	2 MU	4	2	4	2	0	0	0	
	TH.	9622	9750	9845	10064	10080	10085	10233	
	EXP.	9622							
1D 2 (16967)	2 MU	4	0	4	2				
	TH.	16669	16820	16907	17205				
	EXP.	16654		16890	17237				
3P 0 + 11 6 + 3P 1 (20739, 21260, 21307)	2 MU	0	0	2	4	4	0	2	4
	TH.	20732	20772	20825	20943	21135	21204	21333	21335
	EXP.	20728	20794				21204		
	2 MU	4	2	0	0	2			
	TH.	21342	21358	21613	21635	21709			
	EXP.		21358			21692			
3P 2 (22467)	2 MU	4	4	0	2				
	TH.	22343	22376	22570	22591				
	EXP.	22355	22380	22557	22587				
1S 0 (46900)	2 MU	0							
	TH.	46911							

TABLE 8. ENERGY LEVELS FOR Nd³⁺ IN KY₃F₁₀

LSJ state (centroid)		Energy level							
4I $\frac{7}{2}$ (161)	2 MU	3	1	3	1	1			
	TH.	-11	120	128	195	323			
	EXP.	0	120	131	182	320			
4I $\frac{9}{2}$ (2066)	2 MU	1	3	1	3	1	3		
	TH.	1987	1991	2048	2052	2094	2203		
	EXP.	1979	1986	2049	2053	2104			
4I $\frac{11}{2}$ (4032)	2 MU	1	1	3	3	3	1	3	
	TH.	3935	3958	3959	4025	4065	4076	4193	
	EXP.	3928	3967	3956		4063	4078	4193	
4I $\frac{13}{2}$ (6050)	2 MU	1	1	3	3	3	1	3	1
	TH.	5894	5903	5935	5983	6116	6157	6172	6248
	EXP.	5886			5988		6165	6165	6250
4F $\frac{3}{2}$ (11512)	2 MU	3	1						
	TH.	11454	11543						
	EXP.	11450	11547						
4F $\frac{5}{2}$ + 2H $\frac{5}{2}$ (12480, 12590)	2 MU	3	1	1	3	1	1	3	3
	TH.	12394	12475	12523	12524	12543	12621	12624	12673
4F $\frac{7}{2}$ + 4S $\frac{3}{2}$ (13500, 13500)	2 MU	1	3	1	3	3	1		
	TH.	13405	13491	13494	13513	13533	13565		
4F $\frac{9}{2}$ (14700)	2 MU	1	3	1	3	1			
	TH.	14601	14663	14690	14758	14817			
2H $\frac{7}{2}$ (15870)	2 MU	3	1	3	1	3	1		
	TH.	15851	15863	15863	15871	15890	15900		
4G $\frac{7}{2}$ + 2G $\frac{7}{2}$ (17300, 17460)	2 MU	3	1	3	1	1	3	3	
	TH.	17199	17209	17365	17463	17483	17502	17605	

TABLE 9. ENERGY LEVELS FOR Pm^{3+} IN KY_3F_{10}

LSJ state (centroid)		Energy level							
5I 4 (150)	2 MU TH.	0 0	0 11	0 66	2 109	4 179	2 185	4 300	
5I 5 (1577)	2 MU TH.	2 1449	0 1553	4 1565	2 1567	4 1594	0 1594	0 1608	2 1612
5I 6 (3186)	2 MU TH. 2 MU TH.	4 3052 0 3213	4 3053 0 3223	2 3159	0 3167	4 3173	2 3193	4 3202	2 3210
5I 7 (4850)	2 MU TH. 2 MU TH.	2 4703 2 4896	0 4801 0 4904	2 4806 0 4910	4 4807	4 4814	4 4821	4 4883	2 4886
5I 8 (6600)	2 MU TH. 2 MU TH.	0 6433 4 6668	0 6434 2 6694	0 6475 2 6711	2 6479 0 6721	2 6503 0 6731	4 6510	4 6622	4 6633
5F 1 (12400)	2 MU TH.	0 12327	2 12398						
5F 2 (12820)	2 MU TH.	2 12729	4 12806	0 12882	4 12887				
5F 3 (13600)	2 MU TH.	4 13490	2 13556	4 13557	2 13652	0 13661			
5S 2 (14300)	2 MU TH.	4 14285	2 14286	4 14287	0 14291				
5F 4 (14650)	2 MU TH.	0 14606	2 14622	4 14629	2 14638	4 14664	0 14666	0 14673	
5F 5 + 3K 6 (15900, 15900)	2 MU TH. 2 MU TH.	0 15762 4 15884 0 16006	2 15795 2 15884 0 16009	4 15854 0 15889	0 15870 2 15894	2 15872 4 15913	0 15872 4 15924	2 15875 4 15928	4 15878 2 15979

TABLE 10. ENERGY LEVELS FOR Sm^{3+} IN KY_3F_{10}

LSJ state (centroid)		Energy level							
$6H \frac{1}{2}$ (36)	2 MU TH.	3 0	1 219	3 273					
$6H \frac{3}{2}$ (1080)	2 MU TH.	1 1107	3 1186	3 1297	1 1299				
$6H \frac{5}{2}$ (2286)	2 MU TH.	1 2320	1 2363	3 2455	3 2500	1 2505			
$6H \frac{7}{2}$ (3608)	2 MU TH.	3 3607	1 3688	1 3771	3 3796	1 3807	3 3831		
$6H \frac{9}{2}$ (5000)	2 MU TH.	3 4919	3 5109	1 5165	3 5172	1 5189	1 5218	3 5220	
$6H \frac{11}{2} +$ $6F \frac{1}{2} +$ $6F \frac{3}{2}$ (6508, 6400, 6630)	2 MU TH. 2 MU TH.	1 6289 1 6765	1 6559 3 6794	1 6633 1 6809	3 6638	3 6659	3 6719	1 6734	3 6762
$6F \frac{5}{2}$ (7100)	2 MU TH.	3 7245	3 7275	1 7296					
$6F \frac{7}{2}$ (8000)	2 MU TH.	3 8131	3 8157	1 8189	1 8199				
$6F \frac{9}{2}$ (9200)	2 MU TH.	1 9339	1 9351	3 9365	3 9367	1 9411			
$6F \frac{11}{2}$ (10500)	2 MU TH.	1 10607	3 10623	3 10664	3 10679	1 10701	1 10723		
$4G \frac{3}{2}$ (17900)	2 MU TH.	3 17907	3 18012	1 18223					
$4F \frac{3}{2}$ (18900)	2 MU TH.	3 19045	1 19071						

TABLE 11. ENERGY LEVELS FOR Eu^{3+} IN KY_3F_{10}

LSJ state (centroid)		Energy level							
7F 0 (27)	2 MU	0							
	TH.	0							
	EXP.	0							
7F 1 (392)	2 MU	0	2						
	TH.	281	408						
	EXP.	278	411						
7F 2 (1067)	2 MU	2	4	0	4				
	TH.	934	1041	1142	1154				
	EXP.	933	1030	1148	1159				
7F 3 (1927)	2 MU	0	2	2	4	4			
	TH.	1858	1890	1914	1988	2019			
	EXP.	1858	1895	1903	2002	2012			
7F 4 (2894)	2 MU	0	0	0	2	4	2	4	
	TH.	2754	2775	2792	2838	2925	3020	3059	
	EXP.	2748	2778	2800	2845		3014	3052	
7F 5 (3941)	2 MU	2	4	0	2	4	2	0	0
	TH.	3742	3926	3931	3940	3975	4005	4021	4064
	EXP.	3739	3929						
7F 6 (4980)	2 MU	2	0	0	4	4	2	4	4
	TH.	4822	4876	4884	4936	4938	5008	5096	5138
	2 MU	2	0						
	TH.	5181	5200						
5D 0 (17277)	2 MU	0							
	TH.	17276							
5D 1 (19031)	2 MU	0	2						
	TH.	19001	19045						
	EXP.	19007	19039						
5D 2 (21500)	2 MU	0	4	4	2				
	TH.	21464	21488	21512	21517				
	EXP.	21458	21478	21514	21530				
5D 3 (24408)	2 MU	4	2	4	2	0			
	TH.	24386	24392	24404	24420	24444			
5L 6 (25400)	2 MU	4	4	0	2	4	4	2	2
	TH.	25204	25209	25242	25266	25333	25414	25486	25566
	2 MU	0	0						
	TH.	25576	25591						

TABLE 12. ENERGY LEVELS FOR Gd³⁺ IN KY₂F₁₀

LSJ state (centroid)		Energy level							
8S $\frac{7}{2}$ (0)	2 MU TH.	1 0	3 0	3 0	1 1				
6P $\frac{7}{2}$ (32200)	2 MU TH.	1 32147	3 32172	3 32213	1 32252				
6P $\frac{5}{2}$ (32780)	2 MU TH.	1 32743	3 32775	3 32806					
6P $\frac{3}{2}$ (33350)	2 MU TH.	3 33331	1 33364						
6I $\frac{7}{2}$ (35930)	2 MU TH.	1 35907	3 35929	1 35931	3 35957				
6I $\frac{5}{2} +$ 6I $\frac{3}{2}$ (36270, 36340)	2 MU TH. 2 MU TH.	1 36241 3 36339	3 36257 3 36340	1 36269 1 36340	3 36285 1 36342	1 36300 3 36342	3 36339 1 36344	1 36339	1 36339
6I $\frac{5}{2} +$ 6I $\frac{3}{2} +$ 6I $\frac{1}{2}$ (36560, 36660, 36710)	2 MU TH. 2 MU TH. 2 MU TH.	1 36527 3 36648 3 36705	3 36540 3 36657 1 36719	3 36554 1 36673 3 36732	3 36564 1 36687 1 36733	1 36584 3 36687 3 36739	1 36590 1 36690	1 36630 3 36693	3 36637 1 36696
6D $\frac{7}{2}$ (39720)	2 MU TH.	3 39668	1 39675	3 39710	1 39729	1 39816			
6D $\frac{5}{2}$ (40560)	2 MU TH.	1 40550							
6D $\frac{3}{2}$ (40700)	2 MU TH.	1 40689	3 40699	1 40701	3 40707				
6D $\frac{1}{2}$ (40850)	2 MU TH.	1 40836	3 40877						
6D $\frac{3}{2}$ (41000)	2 MU TH.	1 40978	3 40994	3 41057					

TABLE 13. ENERGY LEVELS FOR Tb³⁺ IN KY₃F₁₀

LSJ state (centroid)		Energy level							
7F 6 (85)	2 MU	0	2	4	4	2	4	4	0
	TH.	0	22	71	114	203	288	288	319
	2 MU	0	2						
	TH.	324	384						
7F 5 (2100)	2 MU	0	0	2	4	2	4	0	2
	TH.	2128	2159	2163	2186	2211	2219	2226	2415
7F 4 (3356)	2 MU	4	2	4	2	0	0	0	
	TH.	3342	3373	3448	3526	3575	3591	3602	
7F 3 (4400)	2 MU	4	4	2	2	0			
	TH.	4432	4461	4535	4552	4595			
7F 2 (5038)	2 MU	4	0	4	2				
	TH.	5093	5101	5193	5290				
7F 1 (5440)	2 MU	2	0						
	TH.	5560	5688						
7F 0 (5700)	2 MU	0							
	TH.	5865							
5D 4 (20500)	2 MU	4	2	4	2	0	0	0	
	TH.	20580	20598	20607	20624	20634	20692	20692	
5D 3 + 5G 6 (26336, 26500)	2 MU	2	0	4	4	2	0	2	4
	TH.	26434	26452	26452	26465	26467	26495	26524	26580
	2 MU	4	2	0	0	2	4	4	
	TH.	26601	26665	26672	26673	26678	26744	26745	
5L 10 (27100)	2 MU	2	0	0	4	4	2	0	2
	TH.	27042	27069	27074	27076	27078	27129	27205	27216
	2 MU	0	2	4	4	2	0	4	4
	TH.	27223	27271	27298	27373	27383	27391	27502	27502

TABLE 14. ENERGY LEVELS FOR Dy³⁺ IN KY₃F₁₀

LSJ state (centroid)		Energy level							
6H $\frac{1}{2}$ (52)	2 MU TH.	1 0	3 7	1 31	3 42	3 95	1 97	3 147	1 460
6H $\frac{1}{2}$ (3517)	2 MU TH.	3 3480	1 3502	3 3530	1 3537	1 3554	3 3633	3 3781	
6H $\frac{1}{2}$ (5850)	2 MU TH.	1 5811	3 5828	3 5862	1 5925	1 5991	3 6023		
6H $\frac{3}{2} +$ 6F $\frac{1}{2}$ (7700, 7700)	2 MU TH. 2 MU TH.	3 7638 3 7864	1 7647 1 7875	1 7677 1 7896	3 7698	1 7713	3 7758	3 7792	1 7832
6H $\frac{3}{2} +$ 6F $\frac{3}{2}$ (9100, 9100)	2 MU TH. 2 MU TH.	3 9021 1 9304	1 9057	1 9114	1 9178	3 9180	3 9193	1 9208	3 9250
6H $\frac{3}{2}$ (10200)	2 MU TH.	3 10163	1 10195	3 10435					
6F $\frac{3}{2}$ (11000)	2 MU TH.	1 11042	1 11057	3 11087	3 11133				
6F $\frac{3}{2}$ (12400)	2 MU TH.	1 12445	3 12465	3 12514					
6F $\frac{3}{2}$ (13250)	2 MU TH.	1 13319	3 13336						
6F $\frac{3}{2}$ (13760)	2 MU TH.	1 13839							
4F $\frac{3}{2}$ (21100)	2 MU TH.	3 21035	1 21071	3 21179	1 21194	1 21280			
4I $\frac{1}{2}$ (22100)	2 MU TH.	1 22062	1 22076	3 22101	3 22112	3 22201	1 22214	3 22232	1 22377

TABLE 15. ENERGY LEVELS FOR Ho³⁺ IN KY₃F₁₀

LSJ state (centroid)		Energy level							
5I 8 (108)	2 MU	0	0	2	2	4	4	4	4
	TH.	0	8	18	30	52	82	88	156
	2 MU	2	0	2	0	0			
	TH.	167	182	190	246	256			
5I 7 (5130)	2 MU	0	0	2	2	4	4	2	0
	TH.	5075	5080	5084	5094	5094	5135	5144	5149
	2 MU	4	4	2					
	TH.	5154	5154	5251					
5I 6 (8580)	2 MU	0	0	2	4	2	4	0	2
	TH.	8538	8545	8547	8553	8564	8573	8580	8586
	2 MU	4	4						
	TH.	8692	8692						
5I 5 (11120)	2 MU	2	4	0	4	2	0	0	2
	TH.	11087	11092	11099	11110	11111	11116	11130	11215
5I 4 (13300)	2 MU	4	4	2	2	0	0	0	
	TH.	13175	13258	13261	13310	13333	13409	13417	
5F 5 (15500)	2 MU	0	0	2	4	2	4	2	0
	TH.	15415	15418	15441	15492	15523	15533	15580	15601
5F 4 + 5S 2 (18500, 18500)	2 MU	0	0	4	4	0	2	2	2
	TH.	18434	18439	18477	18481	18481	18507	18527	18534
	2 MU	4	4	0					
	TH.	18538	18545	18558					

TABLE 16. ENERGY LEVELS FOR Er³⁺ IN KY₃F₁₀

LSJ state (centroid)		Energy level							
		1	3	1	3	3	3	1	1
4I $\frac{1}{2}$ (108)	2 MU TH.	0	72	82	109	194	233	261	269
4I $\frac{3}{2}$ (6600)	2 MU TH.	3 6517	1 6618	3 6620	3 6637	1 6693	3 6704	1 6719	
4I $\frac{5}{2}$ (10250)	2 MU TH.	3 10221	1 10290	1 10296	3 10306	1 10325	3 10334		
4I $\frac{7}{2}$ (12400)	2 MU TH.	1 12366	1 12424	3 12445	1 12451	3 12538			
4F $\frac{7}{2}$ (15250)	2 MU TH.	1 15217	3 15251	3 15314	1 15319	1 15378			
4S $\frac{3}{2}$ (18350)	2 MU TH.	1 18346	3 18423						
2H $\frac{1}{2}$ (19150)	2 MU TH.	1 19143	1 19163	3 19190	3 19209	1 19244	3 19250		
4F $\frac{1}{2}$ (20450)	2 MU TH.	1 20444	3 20476	3 20481	1 20588				
4F $\frac{3}{2}$ (22100)	2 MU TH.	3 22127	1 22127	3 22179					
4F $\frac{5}{2}$ (22500)	2 MU TH.	1 22518	3 22599						
2G $\frac{3}{2}$ (24550)	2 MU TH.	1 24518	3 24586	1 24594	1 24608	3 24670			
4G $\frac{1}{2}$ (26400)	2 MU TH.	1 26379	1 26389	3 26419	3 26474	1 26505	3 26547		

TABLE 17. ENERGY LEVELS FOR Tm³⁺ IN KY₃F₁₀

LSJ state (centroid)		Energy level							
3H 6 (170)	2 MU TH. 2 MU TH.	4 0 0 446	4 0 0 465	2 333	0 353	4 389	2 409	4 420	2 439
3F 4 (5900)	2 MU TH.	0 5940	2 6006	0 6015	0 6016	4 6101	2 6191	4 6210	
3H 5 (8400)	2 MU TH.	2 8309	0 8568	0 8603	2 8610	4 8628	0 8643	4 8669	2 8676
3H 4 (12700)	2 MU TH.	0 12708	0 12726	2 12882	4 12896	4 12909	0 12912	2 12956	
3F 3 (14500)	2 MU TH.	0 14615	2 14670	4 14674	4 14701	2 14722			
3F 2 (15100)	2 MU TH.	2 15217	0 15283	4 15340	4 15378				
1G 4 (21350)	2 MU TH.	0 21337	0 21338	0 21378	2 21474	4 21601	2 21678	4 21753	
1D 2 (28000)	2 MU TH.	0 28078	2 28147	4 28252	4 28266				
1I 6 (34900)	2 MU TH. 2 MU TH.	2 34734 2 35436	0 34851 0 35486	0 34858	4 34893	4 34893	2 35063	4 35234	4 35328
3P 0 (35500)	2 MU TH.	0 35677							
3P 1 (36400)	2 MU TH.	2 36518	0 36697						
3P 2 (38250)	2 MU TH.	2 38285	0 38441	4 38534	4 38628				
1S 0 (79590)	2 MU TH.	0 79769							

The phenomenological B_{kq} for Pr³⁺ in KY₃F₁₀ and the odd-k A_{kq} given in table 2 (for an F charge of $q = -1$) were used along with the radial integrals given elsewhere⁶ to calculate electric and magnetic dipole transition probabilities between the Stark split energy levels for Pr³⁺. Branching ratios and lifetimes also were determined. The salient features of these calculations are given in tables 19 and 20. In table 19, the branching ratios from the fluorescing ³P₀ multiplet to the

⁶ Clyde A. Morrison, Nick Karayianis, and Donald E. Wortman, Rare-Earth Ion-Host Lattice Interactions 4. Predicting Spectra and Intensities of Lanthanides in Crystals, Harry Diamond Laboratories HDL-TR-1816 (June 1977).

TABLE 18. ENERGY LEVELS FOR Yb³⁺ IN KY₃F₁₀

LSJ state (centroid)		Energy level			
2F $\frac{7}{2}$ (250)	2 MU TH.	3 0	1 126	3 299	1 490
2F $\frac{5}{2}$ (10450)	2 MU TH.	3 10266	3 10361	1 10664	

TABLE 19. BRANCHING RATIOS FROM ³P₀ ENERGY LEVEL TO LOWER LYING J-MULTIPLETS FOR Pr³⁺ IN KY₃F₁₀ AND IN YLF

Branching ratio	³ P ₀ to (J-multiplet)	KY ₃ F ₁₀	YLF
β_1	(³ H ₄)	0.33	0.36
β_2	(³ H ₅)	0.042	0.041
β_3	(³ H ₆)	0.32	0.50
β_4	(³ F ₂)	0.19	0.021
β_5	(³ F ₃)	0.0053	0.0056
β_6	(³ F ₄)	0.057	0.057
β_7	(¹ G ₄)	0.053	0.022
β_8	(¹ D ₂)	0.00033	0.00003

TABLE 20. ELECTRIC DIPOLE TRANSITION PROBABILITIES FROM FLUORESCING ³P₀ ENERGY LEVEL TO ³H₄ ENERGY LEVELS FOR Pr³⁺ IN KY₃F₁₀ AND IN YLF

KY ₃ F ₁₀ ^a				YLF ^b			
I ^c	2	3	4	I ^c	2	3	4
2.168 × 10 ⁻⁵	σ	E	0	9.674 × 10 ⁻⁶	π	Γ_2	0
0	—	B ₁ or B ₂	70 ^d	1.600 × 10 ⁻⁵	σ	$\Gamma_{3,4}$	79
3.813 × 10 ⁻⁶	π	A ₁ or A ₂	166	—	—	Γ_1	217 ^d
0	—	B ₁ or B ₂	175	—	—	Γ_1	220
5.851 × 10 ⁻⁶	σ	E	217	6.183 × 10 ⁻⁸	σ	$\Gamma_{3,4}$	496
1.585 × 10 ⁻¹²	π	A ₁ or A ₂	510 ^d	—	—	Γ_1	512 ^d
1.196 × 10 ⁻⁵	π	A ₁ or A ₂	540 ^d	8.458 × 10 ⁻⁷	π	Γ_2	514 ^d

^a The ³P₀ is denoted A₁ and is at 20,728 cm⁻¹ in KY₃F₁₀.^b The ³P₀ is denoted Γ_1 and is at 20,866 cm⁻¹ in YLF.^c The most intense ³P₀ to ³H₄ transition was measured to go to the ground state for Pr³⁺ in KY₃F₁₀ and in YLF. Measurements show that the next most intense transitions go to a 79-cm⁻¹ level in YLF and to the 166- and 217-cm⁻¹ levels in KY₃F₁₀.^d Calculated value.

Notes:

1: Electric dipole transition probability from ³P₀.

2: Polarization of emitted radiation.

3: Final state group property designation; the point symmetry at the impurity ion site is C_{4v} in KY₃F₁₀ and is S₄ in YLF.4: Final state energy position in units cm⁻¹.

lower lying J-multiplets are given. Corresponding quantities for Pr^{3+} in YLF are shown for comparison. Similar values of about 0.33 are obtained for β_1 for both materials, which is the first indication that $\text{Pr}^{3+}:\text{KY}_3\text{F}_{10}$ might be as good a blue-green laser as $\text{Pr}^{3+}:\text{YLF}$ would be. In table 20, the $^3\text{H}_4$ energy levels and the individual line-to-line electric dipole transition probabilities are given. These calculations show that the largest transitions from the $^3\text{P}_0$ level in either material would be to the lowest lying energy levels. However, for $\text{Pr}^{3+}:\text{YLF}$, the higher of the possible laser terminal levels is at 79 cm^{-1} ; this occurrence suggests that even at 77 K, $\text{Pr}^{3+}:\text{YLF}$ acts as a three-level system. For Pr^{3+} in KY_3F_{10} , the higher of the possible laser terminal levels is experimentally determined to be at 217 cm^{-1} , and this occurrence might allow four-level laser operation at 77 K. (The Boltzman factor, $e^{-\Delta E/kT}$, which partially determines the population of the terminal level, is 0.14 for the 217 cm^{-1} KY_3F_{10} Pr^{3+} level and is 0.49 for the 79 cm^{-1} YLF level.) On the basis of the above discussions and the results obtained in the above calculations, further work to determine the feasibility of $\text{Pr}^{3+}:\text{KY}_3\text{F}_{10}$ as a blue-green laser seems warranted at this time.

LITERATURE CITED

- (1) L. Esterowitz, R. Allen, M. Kruer, F. Bartoli, L. S. Goldberg, H. P. Jenssen, A. Linz, and V. Nicolai, *J. Appl. Phys.*, **48** (1977), 650.
- (2) J. W. Pierce and H. Y.-P. Hong, *Proceedings of Tenth Rare-Earth Research Conference* (1973), 527.
- (3) Richard P. Leavitt, Clyde A. Morrison, and Donald E. Wortman, *Rare-Earth Ion-Host Crystal Interactions 3. Three-Parameter Theory of Crystal Fields*, Harry Diamond Laboratories HDL-TR-1673 (June 1975).
- (4) P. Porcher and P. Caro, *J. Chem. Phys.*, **65** (1976), 89.
- (5) N. Karayianis, D. E. Wortman, and H. P. Jenssen, *J. Phys. Chem. Solids*, **37** (1976), 675.
- (6) Clyde A. Morrison, Nick Karayianis, and Donald E. Wortman, *Rare-Earth Ion-Host Lattice Interactions 4. Predicting Spectra and Intensities of Lanthanides in Crystal*, Harry Diamond Laboratories HDL-TR-1816 (June 1977).
- (7) Leon Esterowitz, Filbert J. Bartoli, Roger E. Allen, Donald E. Wortman, Clyde A. Morrison, and Richard P. Leavitt, *Energy Levels and Line Intensities for the Ground Configuration of Pr^{3+} in LiYF_4* , Harry Diamond Laboratories HDL-TR-1875 (February 1979).
- (8) *International Tables*, *I*, Kynoch Press, Birmingham, U.K. (1952), 338, table 225.
- (9) M. Labeau, S. Aleonard, A. Vedrine, R. Boutonnet, and J. C. Cousseins, *Mat. Res. Bull.*, **9** (1974), 615.
- (10) A. Vedrine, R. Boutonnet, R. Sabatier, and J. Cousseins, *Bull. Soc. Chim. Fr.*, **3-4** (1975), 445.
- (11) K. H. Hellwege, *Ann. Physik*, **4** (1948), 95.

ACKNOWLEDGEMENT

We thank Dr. Nick Karayianis of the Harry Diamond Laboratories for suggestions concerning the energy level calculations, Dr. Leon Esterowitz and Dr. Filbert J. Bartoli of the Naval Research Laboratory for their interest in the optical properties of $\text{Pr}^{3+}:\text{KY}_3\text{F}_{10}$, and Dr. Van Nicolai of the Office of Naval Research for partial support of this work. We also thank Dr. A. Linz of the Massachusetts Institute of Technology for the excellent optical quality single crystals of rare-earth doped KY_3F_{10} .

DISTRIBUTION

DEFENSE DOCUMENTATION CENTER
CAMERON STATION, BUILDING 5
ALEXANDRIA, VA 22314
ATTN DDC-TCA (12 COPIES)

COMMANDER
US ARMY RSCH & STD GP (EUR)
BOX 65
FPO NEW YORK 09510
ATTN LTC JAMES M. KENNEDY, JR.
CHIEF, PHYSICS & MATH BRANCH

COMMANDER
US ARMY MATERIEL DEVELOPMENT
& READINESS COMMAND
5001 EISENHOWER AVENUE
ALEXANDRIA, VA 22333
ATTN DRXAM-TL, HQ TECH LIBRARY
ATTN DRCDE, DIR FOR DEV & ENGR
ATTN DRCMD-ST

COMMANDER
US ARMY ARMAMENT MATERIEL
READINESS COMMAND
ROCK ISLAND ARSENAL
ROCK ISLAND, IL 61299
ATTN DRSAR-ASF, FUZE &
MUNITIONS SPT DIV
ATTN DRSAR-LEP-L, TECHNICAL LIBRARY

COMMANDER
US ARMY MISSILE & MUNITIONS
CENTER & SCHOOL
REDSTONE ARSENAL, AL 35809
ATTN ATSK-CTD-F

DIRECTOR
US ARMY MATERIEL SYSTEMS
ANALYSIS ACTIVITY
ABERDEEN PROVING GROUND, MD 21005
ATTN DRXSY-MP

TELEDYNE BROWN ENGINEERING
CUMMINGS RESEARCH PARK
HUNTSVILLE, AL 35807
ATTN DR. MELVIN L. PRICE, MS-44

ENGINEERING SOCIETIES LIBRARY
345 EAST 47TH STREET
NEW YORK, NY 10017
ATTN ACQUISITIONS DEPARTMENT

US ARMY ELECTRONICS TECHNOLOGY
& DEVICES LABORATORY
FORT MONMOUTH, NJ 07703
ATTN DELET-DD

DIRECTOR
DEFENSE ADVANCED RESEARCH PROJECTS AGENCY
ARCHITECT BLDG
1400 WILSON BLVD
ARLINGTON, VA 22209

DIRECTOR
DEFENSE NUCLEAR AGENCY
WASHINGTON, DC 20305
ATTN APTL, TECH LIBRARY

UNDER SECRETARY OF DEFENSE FOR RES AND
ENGINEERING
WASHINGTON, DC 20301
ATTN TECHNICAL LIBRARY (3C128)

OFFICE, CHIEF OF RESEARCH,
DEVELOPMENT, & ACQUISITION
DEPARTMENT OF THE ARMY
WASHINGTON, DC 20310
ATTN DAMA-ARZ-A, CHIEF SCIENTIST
DR. M. E. LASSER
ATTN DAMA-ARZ-B, DR. I. R. HERSHNER

COMMANDER
US ARMY RESEARCH OFFICE (DURHAM)
PO BOX 12211
RESEARCH TRIANGLE PARK, NC 27709
ATTN DR. ROBERT J. LONTZ
ATTN DR. CHARLES BOGOSIAN

COMMANDER
ARMY MATERIALS & MECHANICS RESEARCH
CENTER
WATERTOWN, MA 02172
ATTN DRXMR-TL, TECH LIBRARY BR

COMMANDER
NATICK LABORATORIES
NATICK, MA 01762
ATTN DRXRES-RTL, TECH LIBRARY

COMMANDER
USA FOREIGN SCIENCE & TECHNOLOGY CENTER
FEDERAL OFFICE BUILDING
220 7TH STREET NE
CHARLOTTESVILLE, VA 22901
ATTN DRXST-BS, BASIC SCIENCE DIV

DIRECTOR
USA BALLISTICS RESEARCH LABORATORY
ABERDEEN PROVING GROUND, MD 21005
ATTN DRXBR, DIRECTOR, R. EICHELBERGER
ATTN DRXBR-TB, FRANK J. ALLEN
ATTN DRXBR, TECH LIBRARY
ATTN DRDAR-TSB-S (STINFO)

DISTRIBUTION (Cont'd)

DIRECTOR
ELECTRONIC WARFARE LABORATORY
FT MONMOUTH, NJ 07703
ATTN TECHNICAL LIBRARY
ATTN J. CHARLTON
ATTN DR. HIESLMAIR
ATTN J. STROZYK
ATTN DR. E. J. TEBO

DIRECTOR
NIGHT VISION & ELECTRO-OPTICS LABORATORY
FT BELVOIR, VA 22060
ATTN TECHNICAL LIBRARY
ATTN R. BUSER

COMMANDER
ATMOSPHERIC SCIENCES LABORATORY
WHITE SANDS MISSILE RANGE, NM 88002
ATTN TECHNICAL LIBRARY

DIRECTOR
DEFENSE COMMUNICATIONS ENGINEER CENTER
1860 WIEHLE AVE
RESTON, VA 22090
ATTN PETER A. VENA

COMMANDER
US ARMY MISSILE RESEARCH
& DEVELOPMENT COMMAND
REDSTONE ARSENAL, AL 35809
ATTN DRDMI-TB, REDSTONE SCI
INFO CENTER
ATTN DRCPM-HEL, DR. W. B. JENNINGS
ATTN DR. J. P. HALLOWES
ATTN T. HONEYCUTT

COMMANDER
EDGEWOOD ARSENAL
EDGEWOOD ARSENAL, MD 21010
ATTN SAREA-TS-L, TECH LIBRARY

COMMANDER
US ARMY ARMAMENT RES & DEV COMMAND
DOVER, NJ 07801
ATTN DRDAR-TSS, STINFO DIV

COMMANDER
US ARMY TEST & EVALUATION COMMAND
ABERDEEN PROVING GROUND, MD 21005
ATTN TECH LIBRARY

COMMANDER
US ARMY ABERDEEN PROVING GROUND
ABERDEEN PROVING GROUND, MD 21005
ATTN STEAP-TL, TECH LIBRARY, BLDG 305

COMMANDER
WHITE SANDS MISSILE RANGE, NM 88002
ATTN DRSEL-WL-MS, ROBERT NELSON

COMMANDER
GENERAL THOMAS J. RODMAN LABORATORY
ROCK ISLAND ARSENAL
ROCK ISLAND, IL 61201
ATTN SWERR-PL, TECH LIBRARY

COMMANDER
USA CHEMICAL CENTER & SCHOOL
FORT MCCLELLAN, AL 36201

COMMANDER
NAVAL OCEAN SYSTEMS CENTER
SAN DIEGO, CA 92152
ATTN TECH LIBRARY

COMMANDER
NAVAL SUPFACE WEAPONS CENTER
WHITE OAK, MD 20910
ATTN WX-40, TECHNICAL LIBRARY

DIRECTOR
NAVAL RESEARCH LABORATORY
WASHINGTON, DC 20390
ATTN CODE 2620, TECH LIBRARY BR
ATTN CODE 5554, DR. LEON ESTEROWITZ
ATTN CODE 5554, DR. S. BARTOLI
ATTN CODE 5554, R. E. ALLEN

COMMANDER
NAVAL WEAPONS CENTER
CHINA LAKE, CA 93555
ATTN CODE 753, LIBRARY DIV

OFFICE OF NAVAL RESEARCH
ARLINGTON, VA 22217
ATTN DR. V. O. NICOLAI

COMMANDER
AF ELECTRONICS SYSTEMS DIV
L. G. HANSCOM AFB, MA 01730
ATTN TECH LIBRARY

DEPARTMENT OF COMMERCE
NATIONAL BUREAU OF STANDARDS
WASHINGTON, DC 20234
ATTN LIBRARY
ATTN DR. W. BROWNER
ATTN H. S. PARKER

DIRECTOR
LAWRENCE RADIATION LABORATORY
LIVERMORE, CA 94550
ATTN DR. MARVIN J. WEBER
ATTN DR. HELMUT A. KOEHLER

NASA GODDARD SPACE FLIGHT CENTER
GREENBELT, MD 20771
ATTN CODE 252, DOC SECT, LIBRARY

DISTRIBUTION (Cont'd)

NATIONAL OCEANIC & ATMOSPHERIC ADM
ENVIRONMENTAL RESEARCH LABORATORIES
BOULDER, CO 80302
ATTN LIBRARY, R-51, TECH REPORTS

CARNEGIE MELLON UNIVERSITY
SCHENLEY PARK
PITTSBURGH, PA 15213
ATTN PHYSICS & EE
DR. J. O. ARTMAN

UNIVERSITY OF MICHIGAN
COLLEGE OF ENGINEERING NORTH CAMPUS
DEPARTMENT OF NUCLEAR ENGINEERING
ANN ARBOR, MI 48104
ATTN DR. CHIHIRO KIKUCHI

DIRECTOR
ADVISORY GROUP ON ELECTRON DEVICES
201 VARICK STREET
NEW YORK, NY 10013
ATTN SECTRY, WORKING GROUP D

CRYSTAL PHYSICS LABORATORY
MASSACHUSETTS INSTITUTE OF TECHNOLOGY
CAMBRIDGE, MA 02139
ATTN DR. A. LINZ
ATTN DR. H. P. JENSSEN (10 COPIES)

CENTER FOR LASER STUDIES
UNIVERSITY OF SOUTHERN CALIFORNIA
LOS ANGELES, CA 90007
ATTN DR. L. G. DE SHAZER

JOHNS HOPKINS UNIVERSITY
PHYSICS DEPARTMENT
BALTIMORE, MD 21218
ATTN PROF B. R. JUDD

ARGONNE NATIONAL LABORATORY
9700 SOUTH CASS AVENUE
ARGONNE, IL 60439
ATTN DR. W. T. CARNALL

MONTANA STATE UNIVERSITY
PHYSICS DEPT
BOZEMAN, MT 59715
ATTN PROF RUFUS L. CONE

NORTH DAKOTA STATE UNIVERSITY
PHYSICS DEPT
FARGO, ND 58102
ATTN PROF J. GRUBER

RARE-EARTH INFORMATION CENTER
ENERGY AND MINERAL RESOURCES RESEARCH INSTITUTE
IOWA STATE UNIVERSITY
AMES, IA 50011
ATTN DR. KARL GSCHNEIDNER, JR.

US ARMY ELECTRONICS RESEARCH
& DEVELOPMENT COMMAND
ATTN TECHNICAL DIRECTOR, DRDEL-CT

HARRY DIAMOND LABORATORIES
ATTN 00100, COMMANDER/TECH DIR/TSO
ATTN CHIEF, DIV 10000
ATTN CHIEF, DIV 20000
ATTN CHIEF, DIV 30000
ATTN CHIEF, DIV 40000
ATTN RECORD COPY, 81200
ATTN HDL LIBRARY, (3 COPIES) 81100
ATTN HDL LIBRARY, (WOODBIDGE) 81100
ATTN TECHNICAL REPORTS BRANCH, 81300
ATTN CHAIRMAN, EDITORIAL COMMITTEE
ATTN FARRAR, R., 13500
ATTN GLEASON, T., 15400
ATTN KARAYIANIS, N., 13200
ATTN KULPA, S., 13200
ATTN LEAVITT, R., 13200 (10 COPIES)
ATTN MORRISON, C., 13200 (10 COPIES)
ATTN NEMARICH, J., 13300
ATTN WORTMAN, D., 13200
ATTN SATTLER, J., 13200
ATTN SIMONIS, G., 13200
ATTN WORCHESKY, T., 13200